



THE PRACTITIONER LE PRATICIEN

The occasional breech

Peter Hutten-Czapski,
MD

Haileybury, Ont.

Assistant Professor,
Northern Ontario School
of Medicine

Alistair Anderson, MD

Bella Coola, BC

Clinical Assistant Professor,
University of British
Columbia, Vancouver, BC

Correspondence to:
Dr. Peter Hutten-Czapski,
295 Meridian Ave., PO Bag
5000, Haileybury ON
P0J 1K0

This article has been peer
reviewed.

INTRODUCTION

About 3% of pregnancies at term are breech presentations.¹ Some of the perinatal morbidity associated with a breech may be mitigated by planned cesarean section (C-section), at the cost of potential maternal surgical complications.² Pre-emptive C-section for breech presentations has become commonplace. However, even in city hospitals not all women make it to the operating room in time. Thus it is certain that rural doctors, some without local C-section capacity,³ will continue to have to deliver babies who present by the buttocks or feet. It is hoped that this primer will be of assistance in such an occurrence.

[If you are in a hurry, skip to the parts highlighted in bold.]

Breech is associated with increased morbidity. Part of this relates to factors beyond direct control, such as prematurity and congenital abnormalities. Part relates to the challenge of delivering the largest part of the baby, its head.

Over the last 20 years, vaginal breech delivery in North America has all but vanished and has been replaced with the comfortable C-section. The practitioners who retained competency, with historic volume supplemented with a few highly selected perfect breeches (term, frank, multip, generous pelvis and a touch-small fetus), have been mostly convinced by either the lawyers or the Term Breech Trial⁴ to deliver by a booked C-section. Although the results of the Term Breech Trial have not held in the 2-year follow-up⁵ and there are still arguments for planned vaginal delivery,^{6,7} more than 90% of hospitals in one survey delivered breech babies by elective C-section.⁸

So it is not without some trepidation that a rural doctor contemplates the discovery of breech presentation in labour. As labour progresses, the neonatal advantage of C-section is lost.⁹ A rural doctor in such a situation has to contemplate vaginal delivery. Ultimately, what historically made a good breech to deliver, is rendered moot if the baby's sacrum is crowning.

DELIVERING A BREECH

The good news is that the vast majority of breech babies, and hopefully especially those in a hurry, will deliver themselves. Thus, after **calling for help**, the first step in vaginal breech delivery is to encourage the mother's efforts. Both the extra hands and a good rapport will reduce both the mother's and your anxiety and will help you when you do need to intervene.

As the breech crowns, consider a generous episiotomy. It won't help increase room in the pelvis but will make it easier for you to place your hands or forceps. **Do not apply traction on the baby because this may deflect and trap the head and cause injury** (Fig. 1).

When the umbilicus delivers you may pull out a length of cord to ensure slack and to monitor fetal pulse. **If the baby is facing up (sacrum posterior), rotate the baby gently by two hands on its pelvis so it assumes the more favourable face down position.** Allow the baby's legs to deliver by "popping out."

At this point the baby can be either left to hang or supported at 45 degrees to the floor or on a horizontal angle. Do not elevate the body beyond the horizontal (Fig. 2). The baby's back can be

rotated from one anterior oblique to the other, which is helpful in flexing the arms across the chest. The shoulders can be delivered with the trunk in the oblique. When the scapulae deliver, the arms can be optionally swept across the chest and out of the birth canal (Fig. 3, Fig. 4).

Delivery of the head should be by flexion, which presents the same favourable diameters to delivery as with a vertex presentation. A modified Mauriceau–Smellie–Veit (MSV) manoeuvre is used to flex the head. To deliver the head, set yourself below the baby. One hand goes on the baby's back with a finger pushing down on the occiput. Place the other hand under the baby with the forearm supporting it and with two fingers pushing up on

the maxillae. Your assistant will follow with transabdominal pressure flexing the occiput. Some traction on the shoulders by your upper hand may be required. As the head delivers keep the baby's body in neutral position in respect to the head by raising it gently in a large arc (Fig. 5).

The vast majority of babies presenting in the breech position will be delivered by this method.

COMPLICATIONS: PIPER FORCEPS

Forceps delivery is to be respected, as the instrument will amplify leverage and traction, greatly increasing the chance of both delivery and injury. Routine forceps have been advocated for pre-term




Fig. 1. Axial traction is to be avoided as it may cause the head to deflect and result in spinal injury. Keep the sacrum anterior.




Fig. 3. When the scapulae deliver you can assist delivery of the arms by sweeping them across the chest and then extending the elbows.




Fig. 2. The trunk is supported in the horizontal, with a small loop of cord pulled out. A towel can be used to support the trunk in the horizontal plane.




Fig. 4. Gentle rotation of the shoulder girdle facilitates delivery of the left arm.

breech but should be avoided by the inexperienced physician because vaginal breech delivery normally has a good outcome. Early recognition of abnormal breech presentation and attempted delivery is important. **Failure to manually deliver the head in 2 or 3 minutes is an emergency and warrants an attempt at forceps,** by an informed, even if inexperienced, operator.

An assistant holds the baby up to ease application. The operator starts by test assembling the forceps (Pipers are preferred, but any will do) and visualizing the application as if the presentation was occiput anterior. **The handle of the left blade is held by the operator's left hand and inserted almost horizontally into the mother's left side.** The operator's right hand may be used against the patient's left vaginal wall to direct the blade and reduce chances of injury from the insertion (Fig. 6). The blade may be left there or supported by the assistant while the right blade is applied.

The handle of the right blade is held by the operator's right hand and inserted in a manner similar to the first blade, between the mother's right side and the baby's head. The operator's left hand may be used against the patient's right vaginal wall to direct the blade and reduce chances of injury from the insertion.

The handles of the forceps should come together and lock easily without undue force. If not, they should be removed and reapplied. There is no other check on the application that is possible. When the operator is satisfied, **the baby is laid down on the handle of the forceps and traction is applied. At first, traction is applied downward and then, as the head descends, the forceps can be progres-**




Fig. 5. Maintain cephalic flexion by pressure on the baby's maxilla (not mandible!). Delivery of the head is easily accomplished with continued expulsive forces from above and gentle downward traction. As the delivery proceeds, the baby's position moves from the horizontal to a more vertical one.

sively lifted in an arc reflecting the pelvic curve (Fig. 7).

COMPLICATIONS: NUCHAL ARM

One or more of the baby's arms may extend past the neck along with the head. With a generous pelvis or a small baby this may not even be noticed and the delivery effected regardless. However, an attempt to flex the arm across the baby's chest and out of the vagina should be made.

COMPLICATIONS: CERVICAL ENTRAPMENT

Particularly in premature or footling breeches, the body can deliver without dilating the cervix enough




Fig. 6. The baby is held up by the assistant as Piper forceps are applied.




Fig. 7. Traction on the forceps is applied initially below the horizontal plane but rotates through the horizontal toward the vertical as the head delivers.

to allow passage of the after-coming head. At some risk to the mother, pairs of ring forceps can be placed at 2:00, 10:00 and 6:00 o'clock away from the cervical arteries. Radial (Dührssen's) incisions can be made, extending about 3 cm between each pair of forceps to release the head. Adequate exposure, anaesthesia and hemostasis are significant problems, so this should be considered a method of last resort.

CONCLUSIONS

Most, if not all, breech deliveries should be done by elective C-section. A rural doctor, forced by circumstances to deliver a breech baby vaginally, can, in the vast majority of cases, deliver the baby without incident. Very rarely, emergency application of forceps will be required.

Competing interests: None declared.

REFERENCES

1. Hickok DE, Gordon DC, Milberg JA, Williams MA, Daling JR. The frequency of breech presentation by gestational age at birth: a large population-based study. *Am J Obstet Gynecol* 1992;166: 851-2.
2. Hofmeyr GJ, Hannah ME. Planned caesarean section for term breech delivery. *Cochrane Database Syst Rev* 2003;3:CD000166.
3. Iglesias S, Grzybowski SCW, Klein MC, Gagné GP, Lalonde A. Rural obstetrics. Joint position paper on rural maternity care. *Can J Rural Med* 1998;3(2):75-80. Also available in *Can Fam Physician* 1998;44:831-6 and *J Soc Obstet Gynaecol Can* 1998;20(4):393-8.
4. Hannah ME, Hannah W, Hewson SA, Hodnett ED, Saigal S, Willan AR. Planned caesarean section versus planned vaginal birth for breech presentation at term: a randomised multicentre trial. Term Breech Trial Collaborative Group. *Lancet* 2000;356:1375-83.
5. Whyte H, Hannah ME, Saigal S, Hannah WJ, Hewson S, Amankwah K, et al; Term Breech Trial Collaborative Group. Outcomes of children at 2 years after planned cesarean birth versus planned vaginal birth for breech presentation at term: the International Randomized Term Breech Trial. *Am J Obstet Gynecol* 2004;191:864-71.
6. Ulander VM, Gissler M, Nuutila M, Ylikorkala O. Are health expectations of term breech infants unrealistically high? *Acta Obstet Gynecol Scand* 2004;83(2):180-6.
7. Kotaska A. Inappropriate use of randomised trials to evaluate complex phenomena: case study of vaginal breech delivery. *BMJ* 2004;329:1039-42.
8. Hogle KL, Kilburn L, Hewson S, Gafni A, Wall R, Hannah ME. Impact of the International Term Breech Trial on clinical practice and concerns: a survey of centre collaborators. *J Obstet Gynaecol Can* 2003;25(1):14-6.
9. Su M, McLeod L, Ross S, Willan A, Hannah WJ, Hutton E, et al; Term Breech Trial Collaborative Group. Factors associated with adverse perinatal outcome in the Term Breech Trial. *Am J Obstet Gynecol* 2003;189(3):740-5.

DIRECTIVES AUX AUTEURS

Le *Journal canadien de la médecine rurale (JCMR)* est un trimestriel critiqué par les pairs disponible sur papier et sur Internet. Le *JCMR* vise à promouvoir la recherche sur les questions de santé rurale, à promouvoir la santé des communautés rurales (y compris les communautés autochtones), à appuyer et informer les praticiens en milieu rural, à offrir une tribune de débat et de discussion sur la médecine rurale, ainsi qu'à fournir de l'information clinique pratique aux praticiens en milieu rural et à agir sur la politique de santé rurale en publiant des articles qui éclairent les décideurs.

On étudiera la possibilité de publier des documents sur ces questions.

Articles originaux : études de recherche, rapports de cas et analyses critiques d'écrits en médecine rurale. **Commentaires :** éditoriaux, analyses régionales, articles d'opinion. **Articles cliniques :** articles pratiques portant sur la pratique en milieu rural. On encourage la présentation d'illustrations et de photos. **Détexte :** pot-pourri de documents d'intérêt général pour les médecins ruraux (p. ex., voyages, réflexions sur la vie rurale, dissertations). **Couverture :** (Œuvre d'art à thème rural.

Présentation des manuscrits : Veuillez envoyer trois copies papier du manuscrit à l'adresse suivante : Rédacteur en chef, *JCMR*, CP 1086, Shawville (Qué.) J0X 2Y0; 819 647-2972; télécopieur : 819 647-9972; présentez également une version électronique à cjrm@lino.com. Il faut joindre une lettre d'accompagnement indiquant que le texte n'a pas été publié ni

soumis pour publication ailleurs. Les copies papier du manuscrit doivent être dactylographiées à double interligne, comporter une page titre distincte, un résumé d'au plus 200 mots, suivi du texte, des références complètes et des tableaux (chaque tableau sur une page distincte).

«Exigences uniformes pour les manuscrits présentés aux revues biomédicales» : voir www.cmaj.ca/misc/fora-fr.shtml

Permissions écrites : Il faut produire une autorisation écrite des personnes concernées pour utiliser des documents déjà publiés ou des illustrations identifiant des sujets humains, ainsi que de toute personne mentionnée dans les remerciements ou citée comme source d'une communication personnelle.

Illustrations et figures électroniques : Les illustrations doivent être sous forme de clichés de bonne qualité, non montés, sur papier luisant d'au plus 8 sur 10 po (20,3 × 25,4 cm). Les figures présentées sur support électronique doivent être conformes aux caractéristiques techniques décrites à www.amc.ca/cjrm (cliquez sur «Directives aux auteurs»).

Références : Veuillez vous assurer que les références des documents électroniques ne sont pas préparées à l'aide des fonctions notes en fin de texte ou notes en pied de page des logiciels.

Manuscrits acceptés : Les auteurs devront présenter la version la plus récente du manuscrit par courriel ou sur disquette. Veuillez indiquer le logiciel utilisé.